arXiv:1212.2796v1 [math.DG] 12 Dec 2012

NEW SURFACES WITH GENUS ONE IN $\mathbb{H}^2 \times \mathbb{R}$

JULIA PLEHNERT

Abstract. We construct new constant mean curvature surfaces in $\mathbb{H}^2 \times \mathbb{R}$. They arise as sister surfaces of Plateau solutions. It is a family of MC $1/2$ surfaces with k ends, genus 1 and k-fold dihedral symmetry, $k \geq 3$. The surfaces are Alexandrov-embedded.

1. Introduction

For minimal surfaces there exist different approaches to define them, for example as solution of Plateau's problem. In 1970 Lawson proved a correspondence between isometric surfaces in spaceforms: For example, each H-surface (mean curvature $H \equiv 1$) in $\mathbb{R}^3$ corresponds isometrically to a minimal surface in $\mathbb{S}^3$ such that their Gauß maps coincide and their tangent vectors are rotated by $\pi/2$ (with respect to a proper interpretation of the tangent spaces). Two such surfaces are called conjugate cousins ([Law70], [Gro93], [Kar89]). Daniel proved an equivalent correspondence between surfaces in homogeneous 3-manifolds in [Dan07]. A special case of his theorem yields the so-called sister surfaces: Each simply connected surface with constant mean curvature H in $\Sigma^2(\kappa) \times \mathbb{R}$ corresponds to an isometric minimal surface in the homogeneous 3-manifold $E(\kappa + 4H^2, H)$. Hence, instead of proving the existence of a CMC surface in a product space directly, one solves a Plateau problem in a non-trivial Riemannian fibration with base curvature $\kappa + 4H^2$ and considers its sister.

In the case of periodic CMC surfaces in product spaces the conjugate surface construction may be outlined as follows: Consider a geodesic polygon the auxiliary 3-manifold $E(\kappa + 4H^2, H)$, which consists of vertical and horizontal edges. Solve the Plateau problem for the given curve. In the case of unbounded fundamental domains we have to take a limit of the sequence of minimal surfaces. Afterwards solve the period problem(s), i.e. choose the parameters of the geodesic polygon such that the resulting CMC sister surface has the desired properties. With those parameters we constructed a minimal disk whose sister surface is a fundamental domain of the CMC surface. We use Schwarz reflection to establish a complete smooth surface in a product space. In the last step we prove geometric properties of the surface, such as (Alexandrov-)embeddedness, the behaviour of the ends, the genus, etc.

One challenge in the construction in homogeneous manifolds is, that the normals coincide up to their vertical projection in the sense of a Riemannian fibration only. But there are more problems that occur in the construction. In order to prove the existence of periodic surfaces via conjugate construction, the Jordan curve that bounds the Plateau solution of disk-type has to consists of geodesics.

Date: December 13, 2012.
2010 *Mathematics Subject Classification.* Primary 53A10.

The Daniel correspondence implies that a geodesic in the boundary of a minimal surface corresponds to a curvature line in the boundary of the isometric CMC surface, therefore Schwarz reflection applies and continues the surface smoothly, see [GK10] and [MT11]. In the case of conjugate minimal surfaces in $\mathbb{R}^3$ the total curvature of the curvature line in one surface is equal to the total turn of the normal along the corresponding geodesic in the conjugate surface. But in the case of CMC surfaces the total curvature depends also on the length of the curve. In order to solve period problems in horizontal planes, one has to control the total turn of the normal along horizontal curvature lines. For a CMC surface with $H \equiv 1/2$ in $\mathbb{H}^2 \times \mathbb{R}$ this is possible by a horocycle foliation of $\mathbb{H}^2$.

In this paper we prove the existence of new MC 1/2 surfaces in $\mathbb{H}^2 \times \mathbb{R}$, which are k-noids with handle. This work is based on the PhD thesis of the author, [Ple12]. It is organised as follows, we start with an introduction to Riemannian fibration with geodesic fibres and prove a formula for the vertical distance of a horizontal lift of a closed curve.

In Section 3 we cite some properties of Plateau solutions in 3-manifolds and prove, among others, a maximum principle for CMC graphs. In the subsequent section we state some facts about certain sister surfaces and have a closer look on related boundary curves, the sister curves. Section 5 outlines some properties of known minimal and CMC surfaces in Nil_3 and $\mathbb{H}^2 \times \mathbb{R}$, which we use as barriers in our construction.

In the last section we prove the existence of an MC 1/2 surface in $\mathbb{H}^2 \times \mathbb{R}$ with k ends, genus 1 and k-fold dihedral symmetry, $k \geq 3$, which is Alexandrov embedded. We have to solve two period problems in the construction.

Within the last years the theory of minimal in constant mean curvature surfaces in homogeneous 3-manifolds has been evolved actively by many mathematicians. Abresch and Rosenberg introduced a generalized quadratic differential for immersed surfaces in product spaces and show its holomorphicity. Furthermore they classify those surfaces with vanishing differential ([AR05]). Fernandez and Mira constructed a hyperbolic Gauß map to study mean curvature one half surfaces in $\mathbb{H}^2 \times \mathbb{R}$, [FM07]. Recently Cartier and Hauswirth published a work, where they study constant mean curvature 1/2 surfaces in $\mathbb{H}^2 \times \mathbb{R}$ with vertical ends ([CH12]).

2. RIEMANNIAN FIBRATIONS WITH GEODESIC FIBRES

Let us consider a fibration $\pi\colon E \to \Sigma$ of a 3-dimensional complete Riemannian manifold $(E, \langle\cdot,\cdot\rangle)$ over an oriented surface Σ and a vector field ξ tangent to the fibre. We decompose the tangent space $\mathrm{T}_p E$ of the total space E into its horizontal and vertical parts. The *horizontal space* $(\mathrm{T}_p E)^h$ is defined by

$$(\mathrm{T}_p E)^h := \{v \in \mathrm{T}_p E \colon \langle v, \xi(p)\rangle = 0\},$$

the vertical space is given by $(\mathrm{T}_p E)^v := \ker(\mathrm{d}\,\pi_p)$.

We require the fibration to fulfil three properties: It has to be a *Riemannian* fibration with *geodesic fibres* over a 2-dimensional *space form*.

First of all, we consider a *Riemannian fibration*, i.e. $\mathrm{d}\,\pi_p\colon (\mathrm{T}_p E)^h \to \mathrm{T}_{\pi(p)}\Sigma$ is an isometry. This means that the horizontal spaces as well as the tangent space of the base manifold look the same along one fibre. Furthermore, this implies that the horizontal lifts of geodesics in Σ are geodesics in E. For an arbitrary curve γ in Σ

there is always a unique *horizontal lift* $\tilde{\gamma}$ if we fix $\tilde{\gamma}(0) = p \in E$, i.e. $\tilde{\gamma}'(t) \in (\mathrm{T}_{\tilde{\gamma}(t)} E)^h$, for all t.

Secondly, we require the fibres to be geodesics. By computing the Lie derivative, this leads to the following proposition.

Proposition 2.1 ([GK10]). *For a Riemannian fibration $\pi \colon E \to \Sigma$ with geodesic fibres, there exists an unit vector field ξ tangent to the fibres, which is a Killing field.*

We call the vector field ξ the *vertical vector field*. The translations along the fibres are isometries, hence the Killing field ξ generates a subgroup G of $\mathrm{Iso}(E)$. The integral curves of ξ define a principal bundle with connection 1-form $\omega(X) = \langle X, \xi\rangle$. The curvature form is $\Omega := \mathrm{D}\,\omega = \mathrm{d}\,\omega + \frac{1}{2}[\omega,\omega] = \mathrm{d}\,\omega$. The following equation holds for two arbitrary vector fields X, Y

$$\begin{aligned}
\Omega(X,Y) = \mathrm{d}\,\omega(X,Y) &= X\omega(Y) - Y\omega(X) - \omega\left([X,Y]\right) \\
&= \langle \nabla_X Y, \xi\rangle + \langle Y, \nabla_X \xi\rangle - \langle \nabla_Y X, \xi\rangle - \langle X, \nabla_Y \xi\rangle - \left(\langle \nabla_X Y, \xi\rangle - \langle \nabla_Y X, \xi\rangle\right) \\
&= \langle \nabla_X \xi, Y\rangle - \langle \nabla_Y \xi, X\rangle \\
&\overset{(*)}{=} 2\langle \nabla_X \xi, Y\rangle,
\end{aligned}$$

where $(*)$ follows from the fact that ξ is a Killing field. Moreover, the fact that ξ is Killing implies $\xi\Omega(X,Y) = 0$, i.e. Ω is constant along the fibers. Moreover, we claim: For $X^h = X - X^v$ we have $\Omega(X^h, Y^h) = \Omega(X,Y)$. Therefore Ω induces a 2-form $\underline{\Omega} = (\pi^{-1})^*\Omega$ on the base manifold Σ. To see the claim we compute

$$\begin{aligned}
\Omega(X^h, Y^h) &= 2\langle \nabla_{X^h}\xi, Y^h\rangle \\
&= 2\langle \nabla_{(X-X^v)}\xi, Y - Y^v\rangle \\
&= 2(\langle \nabla_X \xi, Y\rangle \underbrace{-\langle \nabla_X \xi, Y^v\rangle}_{=\langle \nabla_{Y^v}\xi, X\rangle} - \langle \nabla_{X^v}\xi, Y\rangle + \langle \nabla_{X^v}\xi, Y^v\rangle) \\
&= 2\langle \nabla_X \xi, Y\rangle,
\end{aligned}$$

because $\nabla_U \xi = 0$ for any vertical vector field U.

Since Σ is oriented, there exists a $\pi/2$-rotation J on $\mathrm{T}_y\,\Sigma$, it induces a $\pi/2$-rotation R on $(\mathrm{T}_p\,E)^h$.

Definition 2.2. Let $\pi \colon E \to \Sigma$ be a Riemannian fibration with geodesic fibres. Its *bundle curvature* τ is a map $\tau \colon \Sigma \to \mathbb{R}$ given by

$$\tau(y) := -\frac{1}{2}\Omega(X, RX) = \frac{1}{2}\langle [X, RX], \xi\rangle,$$

hence $[X, RX]^v = 2\tau(y)\xi$, where X is an arbitrary horizontal unit vector field along $\pi^{-1}(y)$.

We see that Ω measures the non-integrability of the horizontal distribution. Since Ω is constant along the fibres, the map τ is well-defined. The induced 2-form $\underline{\Omega}$ factorizes the natural volume form vol_Σ of Σ

$$-\frac{1}{2}\underline{\Omega} = \tau\,\mathrm{vol}_\Sigma,$$

since we have $\mathrm{vol}_\Sigma(x, Jx) = 1$ and $\underline{\Omega}(x, Jx) = \Omega(\tilde{x}, R\tilde{x})$ for any unit vector x.

The third requirement concerns the base manifold only. If we consider a two-dimensional space form $\Sigma(\kappa)$ as base, the Riemannian fibrations are classified as follows:

Theorem 2.3 ([GK10])**.** *Suppose a Riemannian fibration $\pi\colon E \to \Sigma(\kappa)$ has geodesic fibres, where E is a complete oriented simply connected Riemannian 3-manifold. Then E is a homogeneous manifold with a 4-dimensional isometry group and constant bundle curvature τ.*

We write $E(\kappa,\tau)$ for those spaces. The isometry group of $E(\kappa,\tau)$ depends on the signs of κ and τ, and is equivalent to the isometry group of one of the following Riemannian manifolds:

curv.	$\kappa < 0$	$\kappa = 0$	$\kappa > 0$
$\tau = 0$	$\mathbb{H}^2 \times \mathbb{R}$	$\mathbb{R}^3$	$\mathbb{S}^2 \times \mathbb{R}$
$\tau \neq 0$	$\widetilde{\mathrm{SL}}_2(\mathbb{R})$	$\mathrm{Nil}_3(\mathbb{R})$	(Berger-)$\mathbb{S}^3$

Another interpretation of the bundle curvature is the vertical distance of a horizontal lift of a closed curve:

Lemma 2.4 (Vertical distances)**.** *Let γ be a closed Jordan curve in the base manifold $\Sigma(\kappa)$ of a Riemannian fibration with geodesic fibres $E(\kappa,\tau)$. With Δ defined by $\partial\Delta = \gamma$, we have*

$$d(\tilde{\gamma}(0), \tilde{\gamma}(l)) = 2\tau\,\mathrm{area}(\Delta),$$

where $\tilde{\gamma}$ is the horizontal lift with $\pi(\tilde{\gamma}(0)) = \pi(\tilde{\gamma}(l))$, area *is the oriented volume and $d(p,q)$ denotes the signed vertical distance, which is positive if $\overline{pq}$ is in the fibre-direction ξ.*

Proof. We consider an arclength parametrization $\gamma\colon [0,l] \to \Sigma(\kappa)$ and its horizontal lift $\tilde{\gamma}$. Then $\tilde{\gamma}(0)$ and $\tilde{\gamma}(l)$ are contained in one fibre, i.e. $\pi(\tilde{\gamma}(0)) = \pi(\tilde{\gamma}(l))$. Hence, there exists a vertical arclength parametrized curve c with $c(0) = \tilde{\gamma}(l)$ and $c(v) = \tilde{\gamma}(0)$. The union $c \cup \tilde{\gamma} =: \tilde{\Gamma}$ is a closed curve in $E(\kappa,\tau)$ and c' is parallel to ξ. If $c' = \pm\xi$, then

$$\pm v = \int_0^v \langle c', \xi\rangle = \int_0^v \langle c', \xi\rangle + \int_0^l \langle \tilde{\gamma}', \xi\rangle,$$

since $\tilde{\gamma}'$ is horizontal. By definition of the connection 1-form ω, the sum of the integrals is equal to $\int_{\tilde{\Gamma}} \omega$. We apply Stokes's Theorem to get

$$\int_{\tilde{\Gamma}} \omega = \int_{\tilde{\Delta}} \mathrm{d}\,\omega = \int_{\tilde{\Delta}} \Omega = \int_{\pi(\tilde{\Delta})} (\pi^{-1})^*\Omega,$$

where $\tilde{\Delta}$ is any lift of Δ, such that $\pi\colon \tilde{\Delta} \to \Delta$ is one-to-one and $\partial\tilde{\Delta} = \tilde{\Gamma}$. Since $(\pi^{-1})^*\Omega$ is a 2-form on Σ, it factorizes the natural volume form vol_Σ

$$\int_{\pi(\tilde{\Delta})} (\pi^{-1})^*\Omega = -2\int_{\Delta} \tau\,\mathrm{vol}_\Sigma = -2\tau\,\mathrm{area}(\Delta).$$

We conclude that

$$2\tau\,\mathrm{area}(\Delta) = \begin{cases} -v, & \text{if } c' = \xi, \\ v, & \text{if } c' = -\xi, \end{cases}$$

and $d(\tilde{\gamma}(0), \tilde{\gamma}(l)) = d(c(v), c(0)) = 2\tau\,\mathrm{area}(\Delta)$. □

3. Solution of the Plateau problem

From the 18th century until 1930 it was an open question whether a rectifiable closed curve Γ in $\mathbb{R}^3$ bounds an area minimizing surface. Douglas ([Dou31]) and Radó ([Rad30]) proved independently that there always exists a minimal surface of disc type spanned by Γ. In 1948 Morrey generalised the theorem to minimal surfaces in homogeneously regular Riemannian manifolds without boundary ([Mor66]). By [Oss70] and [Gul73], the least area disc is a minimal immersion in the interior. In the 80ties Meeks and Yau ([MY82]) showed that in manifolds with mean convex boundary the Plateau solution M is even embedded. A Riemannian manifold N with boundary is *mean convex* if the boundary ∂N is piecewise smooth, each smooth subsurface of ∂N has non-negative mean curvature with respect to the inward normal, and there exists a Riemannian manifold N' such that N is isometric to a submanifold of N' and each smooth subsurface S of ∂N extends to a smooth embedded surface S' in N' such that $S' \cap N = S$. We call each surface S a *barrier*.

Under additional assumptions we may exclude boundary branch points. For example, let the Jordan curve $\Gamma \subset E(\kappa, \tau)$ be piecewise geodesic, at each vertex, the angle is of the form π/n, with natural $n \geq 2$, and for each edge there exists an isometry of $E(\kappa, \tau)$ which acts by π rotation about that edge. If the mean convex domain Ω is disjoint under those isometries, then M extends smoothly by Schwarz reflection across Γ. In the case where $\pi(\partial M)$ bounds a proper subset $\Delta \subset \Sigma$ and $M \subset \pi^{-1}(\Delta)$, then M is a section over Δ, see [GK10].

The maximum principle implies uniqueness of constant mean curvature sections with the same boundary data:

Proposition 3.1. *Let $\pi\colon E \to \Sigma$ be a Riemannian fibration with geodesic fibres. Suppose M is a section over $\Delta \subset \Sigma$ with MC H and prescribed boundary values such that $\pi\colon \partial M \to \partial\Delta$ is injective. Then M is unique.*

Proof. Assume that we have two sections M and $\hat{M}$ over Δ with the same boundary values, they fulfil the non-parametric mean curvature equation. Therefore, their parametrizations u and $\hat{u}$ are solutions of the following differential equation:

$$Q(u) := \partial_x\left(\frac{\lambda U}{W}\right) + \partial_y\left(\frac{\lambda V}{W}\right) - 2\lambda^2 H = 0,$$

where $U = u_x + \lambda\tau y$, $V = u_y - \lambda\tau x$, $W = \sqrt{1 + U^2 + V^2}$ and $\lambda = \frac{4}{4+\kappa(x^2+y^2)}$.

This equation is non-linear. We set $a^i(p) := \frac{\lambda p_i}{\sqrt{1+p_1^2+p_2^2}}$ and $R := U\partial_x + V\partial_y$. Considering the difference $u - \hat{u}$ we get:

$$\begin{aligned}
0 = Q(u) - Q(\hat{u}) &= \sum_i \partial_i a^i(R) - \partial_i a^i(\hat{R}) \\
&= \sum_i \partial_i a^i(tR + (1-t)\hat{R})|_{t=0}^{t=1} \\
&= \int_0^1 \frac{\mathrm{d}}{\mathrm{d}\,s}\left[\sum_i \partial_i a^i(sR + (1-s)\hat{R})\right]_{s=t} \mathrm{d}\,t \\
&= \sum_{i,j} \int_0^1 \partial_i \left[\frac{\partial a^i}{\partial p_j}(tR + (1-t)\hat{R})(R_j - \hat{R}_j)\right] \mathrm{d}\,t \\
&= \sum_{i,j} \partial_i \left[\underbrace{\left(\int_0^1 \frac{\partial a^i}{\partial p_j}(tR + (1-t)\hat{R})\,\mathrm{d}\,t\right)}_{=:a^{ij}}(R_j - \hat{R}_j)\right] \\
&= \sum_{i,j} \partial_i [a^{ij}\partial_j(u - \hat{u})].
\end{aligned}$$

Since

$$\partial_j a^i(p) = \frac{\lambda \delta_{ij}}{\sqrt{1+p_1^2+p_2^2}} - \frac{\lambda p_i p_j}{\sqrt{1+p_1^2+p_2^2}^3} = \partial_i a^j(p),$$

then $(a^{ij})_{ij}$ is symmetric, moreover $\left|(\partial_i a^{ij})_j\right|$ is bounded. Therefore with $w := u - \hat{u}$, we have that

$$L(w) := \sum_{i,j} \partial_i(a^{ij}\partial_j w) = Q(u) - Q(\hat{u}) = 0$$

is a linear second order partial differential operator.

With $T(t,x,y) := tR(x,y) + (1-t)\hat{R}(x,y)$ we get $\partial_j a^i(T) = \frac{\lambda\delta_{ij}}{\sqrt{1+|T|^2}} - \frac{\lambda T_i T_j}{\sqrt{1+|T|^2}^3}$. For any compact subset $K \subset \Delta$ the norm $|T|$ has a maximum on $[0,1] \times K$. Hence, there exists $\sigma(K, u, \hat{u}) > 0$, such that we may estimate, by means of the Schwarz inequality,

$$\sum_{i,j} a^{ij}(T)\xi_i\xi_j = \frac{(1+|T|^2)\,|\xi|^2 - \langle T, \xi\rangle^2}{\sqrt{1+|T|^2}^3} \geq \frac{|\xi|^2}{\sqrt{1+|T|^2}^3} > \sigma\,|\xi|^2\,.$$

We conclude that L is uniformly elliptic and therefore, the maximum principle for elliptic partial differential equations ([GT01]) applies. □

Remark 3.2. The uniqueness of a section is also true in a more general case: The projection of ∂M has to be injective except for at most finitely many points of $\partial\Delta$. This means, we allow vertical segments in the boundary. The proof needs a more general maximum principle by Nitsche; for $\mathbb{R}^3$ see [Nit75].

For a sequence of minimal sections there exists a uniform curvature estimate ([GK10]) and a uniform area bound, therefore there exists a limit M if we write them locally as normal graphs over their tangent space.

Proposition 3.3. *Let M_n be a sequence of minimal discs, which are graphs in a Riemannian fibration $\pi\colon E(\kappa,\tau) \to \Sigma(\kappa)$ converging to a minimal disc M. Then M is either a graph over $\lim_{n\to\infty} \pi(M_n)$ or a vertical plane.*

Proof. Since each M_n is a graph, it is transversal to the fibres and for the normal ν_n we have $\langle \nu_n, \xi\rangle > 0$ for each $n \in \mathbb{N}$. For the limit $\nu_n \to \nu$ we still know that $\langle \nu, \xi\rangle \geq 0$. If $\langle \nu, \xi\rangle > 0$, then M is a graph. Otherwise we pick $p \in M$ with $\langle \nu(p), \xi\rangle = 0$ and consider the unique vertical plane $V \subset E(\kappa,\tau)$ that is orthogonal to ν in p. If $M \neq V$, we analyse $\Gamma := M \cap V$. By the maximum principle we have $\Gamma \neq \{p\}$.

Locally in a neighbourhood $U_p \subset V$ of p, the surface M is a graph M_p over V in the normal direction.

We claim: There exists a map $\hat{d}\colon M_p \subset M \to \mathbb{R}$ with $M_p \cap V = \{\hat{d} = 0\}$ such that its composition with a parametrisation f of M is a real analytic function $d := \hat{d} \circ f$.

We construct $\hat{d}$ via the diffeomorphism $e\colon M_p \times (-\delta,\delta) \to E(\kappa,\tau)$ defined by $e(q,t) = \exp^E(t\nu_M(q))$. The map $\hat{d} := \pi_{(-\delta,\delta)} \circ e^{-1}\colon M_p \to \mathbb{R}$ is the signed distance function of M. We have $M \cap U_p = \{\hat{d} = 0\}$.

If we choose a harmonic and conformal parametrisation $f\colon D \to E(\kappa,\tau)$ of M, where $D \subset \mathbb{R}^2$ is the open unit disc, then the function $d := \hat{d} \circ f$ is analytic. Therefore, it can be expressed as a power series $d(x) = \sum_{j=m}^{\infty} P_j(x)$, where P_j is a polynomial of degree j and $P_m \neq 0$. Since the first derivatives of V and M coincide, we have $m \geq 2$.

Moreover, Γ divides U_p in at least $2m$ regions. There exists $q \in M \setminus \Gamma$ with a neighbourhood M_q, such that $M_q \cap \Gamma = \emptyset$ and $\pi_{\Sigma(\kappa)}(M_q)$ is not injective. But since M is the limit of M_n this means that there exists $n \in \mathbb{N}$ for which $\pi_{\Sigma(\kappa)}(M_n)$ is not injective, a contradiction to M_n being a graph.

Therefore, $P_m = 0$ and $d \equiv 0$, which means that M is a vertical plane. □

The fact that the intersection of two non-transversally intersecting minimal surfaces consists of $2m$ embedded curves is true in general (see [CM11]).

4. Sister surfaces

Lawson made a great contribution in the study of constant mean curvature surfaces in 1970 when he showed the correspondence between minimal surfaces in $\mathbb{S}^3$ and H-surfaces in $\mathbb{R}^3$, see [Law70]. By means of the reflection principle in $\mathbb{S}^3$ it was then possible to construct new surfaces in the Euclidean space. In recent years mathematicians have grown their interest within surfaces in other ambient manifolds. In 2007 Daniel published a generalized Lawson correspondence for homogeneous manifolds ([Dan07]), we use one special case of his correspondence:

Theorem 4.1 ([Dan07, Theorem 5.2]). *There exists an isometric correspondence between an MC H-surface $\tilde{M}$ in $\Sigma(\kappa) \times \mathbb{R} = E(\kappa, 0)$ and a minimal surface M in $E(\kappa + 4H^2, H)$. Their shape operators are related by*

$$\tilde{S} = JS + H\,\mathrm{id}, \tag{4.1}$$

where J denotes the $\pi/2$ rotation on the tangent bundle of a surface. Moreover, the normal and tangential projections of the vertical vector fields ξ and $\tilde{\xi}$ are related by

$$\langle\tilde{\xi},\tilde{\nu}\rangle = \langle\xi,\nu\rangle, \qquad J\,\mathrm{d}f^{-1}(T) = \mathrm{d}\tilde{f}^{-1}(\tilde{T}), \tag{4.2}$$

where f and $\tilde{f}$ denote the parametrizations of M and $\tilde{M}$ respectively, ν and $\tilde{\nu}$ their unit normals, and T, $\tilde{T}$ the projections of the vertical vector fields on $\mathrm{T}M$ *and* $\mathrm{T}\tilde{M}$.

We call the isometric surfaces M and $\tilde{M}$ *sister surfaces*, or *sisters* in short.

Examples:

1. For $H = 0$ the surfaces M and $\tilde{M}$ are conjugate minimal surfaces in $\Sigma(\kappa)\times\mathbb{R}$.
2. For $H \in (0,1/2)$ and $\kappa = -1$ we have $4H^2 - 1 < 0$ and therefore corresponds to a minimal surface in $\widetilde{\mathrm{PSL}_2}(\mathbb{R})$ and its MC H-sister surface in $\mathbb{H}^2\times\mathbb{R}$.
3. For $H = 1/2$ and $\kappa = -1$ an MC $1/2$-surface in $\mathbb{H}^2\times\mathbb{R}$ corresponds to a minimal surface in $E(0,1/2) = \mathrm{Nil}_3(\mathbb{R})$.
4. Furthermore, for $H > 1/2$ an MC H-surface in $\mathbb{H}^2\times\mathbb{R}$ results from a minimal surface in the Berger spheres $E(4H^2-1,H)$, since $4H^2-1>0$.

4.1. **Reflection principles and sister curves.** We want to apply Schwarz reflection to construct complete periodic CMC surfaces in $E(\kappa,\tau)$. It is well-known that the Schwarz reflection extends minimal surfaces in space forms (see [Law70]) with respect to the space form symmetries. In $E(\kappa,\tau)$ the isometry group has at least dimension 4, and there are symmetries, for which the Schwarz reflection applies. We reflect across a geodesic $c \subset E(\kappa,\tau)$ or a totally geodesic plane $V \subset E(\kappa,\tau)$, i.e. the geometric interpretation is to send a point p to its opposite point on a geodesic through p that meets c or V orthogonally.

Große-Brauckmann and Kusner proved that if c is a horizontal or vertical geodesic of $E(\kappa,\tau)$, then the geodesic reflection across c is an isometry. Moreover, in the product spaces $E(\kappa,0) = \Sigma(\kappa)\times\mathbb{R}$ a geodesic reflection across vertical or horizontal planes is an isometry ([GK10]). For this isometries we formulate the Schwarz reflection principles: Suppose that a minimal surface is smooth up to the boundary and the boundary contains a curve which is a horizontal or vertical geodesic of $E(\kappa,\tau)$. Then the geodesic is an asymptotic direction and the reflection preserves the principal curvatures, therefore it extends the surface smoothly. A CMC surface in a product space $E(\kappa,0)$, which is smooth up to the boundary, extends smoothly if the boundary contains a curve in a vertical or horizontal plane, provided the surface conormal is perpendicular to the plane, since the curve a curvature direction. A totally geodesic plane is called *mirror plane*, and a curve in which the surface meets a mirror plane orthogonally is called *mirror curve*.

Remark 4.2. We construct surfaces by solving Plateau problems. The solution is an area minimising map from a disc to $E(\kappa,\tau)$, continuous up to the boundary. To apply the Schwarz reflection principles we have to exclude branch points in the interior ([Oss70] and [Gul73]) and at the boundary ([GK10]).

4.2. **Sister curves.** We want to analyse the geometry of periodic surfaces. Therefore, we take a closer look at their boundary curves.

Let $c = f\circ\gamma$ be a curve parametrized by arc length in a hypersurface $M = f(\Omega) \subset \bar{M}$ with (surface) normal ν. The *normal curvature* k and the *normal torsion* t along c are defined by

$$k := \nu\cdot\bar{\nabla}_{c'}c' = -\bar{\nabla}_{c'}\nu\cdot c' = \langle S\gamma',\gamma'\rangle, \qquad t := -\bar{\nabla}_{c'}\nu\cdot Jc' = \langle S\gamma', J\gamma'\rangle.$$

Let $\tilde{M} \subset \Sigma(\kappa) \times \mathbb{R}$ denote an MC H-surface and $M \subset E(\kappa + 4H^2, H)$ its minimal sister. Furthermore, let γ be a curve in Ω. We call $\tilde{c} := \tilde{f}(\gamma)$ and $c := f(\gamma)$ *sister curves*.

Lemma 4.3. *For a pair of sister curves the normal curvature and torsion are related as follows:*

$$\tilde{k} = -t + H \quad \text{and} \quad \tilde{t} = k.$$

Proof. We apply Equation (4.1) to the definitions:

$$\begin{aligned} \tilde{k} &= \langle \tilde{S}\gamma', \gamma' \rangle = \langle (JS + H\,\mathrm{id})\gamma', \gamma' \rangle = -t + H, \\ \tilde{t} &= \langle \tilde{S}\gamma', J\gamma' \rangle = \langle (JS + H\,\mathrm{id})\gamma', J\gamma' \rangle = k. \end{aligned}$$ □

From this follows a relation between mirror curves and their sister curves, see [GK10] and [MT11]:

1. A curve $\tilde{c} \subset \tilde{M} \subset \Sigma(\kappa) \times \mathbb{R}$ is a mirror curve in a vertical plane if and only if its sister curve $c \subset M \subset E(\kappa + 4H^2, H)$ is a horizontal geodesic.
2. Similarly, $\tilde{c}$ is a horizontal mirror curve if and only if c is a vertical geodesic.

In our construction we consider the *fundamental patch* of a periodic MC H-surface. The complete surface is then generated by reflections. The fundamental patch is simply connected and bounded by mirror curves $\tilde{c}_i$ in vertical and horizontal planes. Given a fundamental patch bounded by n arc length parametrized mirror curves $\tilde{c}_i$, it defines the following geometric quantities:

1. The *length* $\tilde{l}_i$ of the mirror curve $\tilde{c}_i$, also denoted by $l(\tilde{c}_i)$ or $|\tilde{c}_i|$.
2. The *vertex angle* $\tilde{\varphi}_i$ of two edges $\tilde{c}_i$ and $\tilde{c}_{i+1}$, which satisfies
$$\cos \tilde{\varphi}_i = -\tilde{c}_i{}'(\tilde{l}_i) \cdot \tilde{c}'_{i+1}(0).$$
3. The total *turn* angle of the normal $\tilde{\nu}$
$$\mathrm{turn}_i = \mathrm{turn}_{\tilde{c}_i}(\tilde{\nu}) := \int_{\tilde{c}_i} \tilde{k},$$
which measures the total turn of the normal relative to a parallel field.

The minimal sister surface is bounded by horizontal and vertical geodesics. Since the surfaces are isometric, we have

$$\tilde{l}_i = l_i, \qquad \tilde{\varphi}_i = \varphi_i. \tag{4.3}$$

Accordingly, we want to measure the rotational angle of the normal ν along a curve c in $E(\kappa + 4H^2, H)$. We detect the *twist* of the normal with respect to an appropiate vector field X

$$\mathrm{twist}_c(\nu, X) := \int_c \langle \nabla_{c'}\nu, c' \times \nu \rangle - \langle \nabla_{c'}X, c' \times X \rangle.$$

Definition 4.4. (1) Let $c \subset M$ be a vertical geodesic in $E(\kappa + 4H^2, H)$, then the twist is defined by the total rotation speed of ν with respect to a basic vector field $\tilde{e}$, i.e. the horizontal lift of any vector field e on $\Sigma(\kappa + 4H^2)$:

$$\mathrm{twist}_v := \mathrm{twist}_c(\nu, \tilde{e}) = \int_c \langle \nabla_{c'}\nu, c' \times \nu \rangle - \langle \nabla_{c'}\tilde{e}, c' \times \tilde{e} \rangle.$$

(2) Let $c \subset M$ be a horizontal geodesic in $E(\kappa+4H^2, H)$, then the twist is defined by the total rotation speed of ν with respect to the vertical vector field ξ:

$$\mathrm{twist}_h := \mathrm{twist}_c(\nu, \xi) = \int\limits_c \langle \nabla_{c'}\nu, c' \times \nu\rangle - \langle \nabla_{c'}\xi, c' \times \xi\rangle.$$

With this definition twist_v measures the angle $\measuredangle(\mathrm{d}\,\pi_c(\nu(0)), \mathrm{d}\,\pi_c(\nu(l))$ in the projection. Please note that the Definition 4.41 differs from the one in [GK10]. We drop the index when it is clear whether the geodesic is vertical or horizontal.

Lemma 4.5. (1) *Let $c \subset M$ be a vertical geodesic in $E(\kappa + 4H^2, H)$, whose sister is a horizontal mirror curve $\tilde{c} \subset \tilde{M}$ in $E(\kappa, 0)$. Then*

$$\mathrm{twist}_v = \int\limits_c t + Hl(c) \quad \textit{and} \quad \tilde{k} = 2H - \mathrm{twist}_v'\,.$$

(2) *Let $c \subset M$ be a horizontal geodesic in $E(\kappa + 4H^2, H)$, whose sister is a vertical mirror curve $\tilde{c} \subset \tilde{M}$ in $E(\kappa, 0)$. Then*

$$\mathrm{twist}_h = -\,\mathrm{turn}\,.$$

Proof. (1) Without loss of generality $c' = \xi$. Let (c', Jc', ν) be positively oriented. Let J and R denote $\pi/2$-rotations in the tangent bundle $\mathrm{T}M$ and the horizontal plane $\mathrm{d}\,\pi^{-1}(\mathrm{T}\,\Sigma(\kappa + 4H^2))$, respectively. Then Jc' and ν are horizontal with $-R\nu = Jc'$.

We denote by E an unit basic vector field. We have

$$\begin{aligned}
\mathrm{twist}_v &= \int\limits_c \left(\langle \nabla_{c'}\nu, c' \times \nu\rangle - \langle \nabla_{c'}E, c' \times E\rangle\right) \\
&= \int\limits_c \left(\langle \nabla_{c'}\nu, R\nu\rangle - \langle \nabla_{c'}E, RE\rangle\right) \\
&= \int\limits_c \left(-\langle \nabla_{c'}\nu, Jc'\rangle - \langle \nabla_{\xi}E, RE\rangle\right) \\
&= \int\limits_c \left(t + H\langle RE, RE\rangle\right) \\
&= \int\limits_c t + Hl(c).
\end{aligned}$$

Lemma 4.3 implies that $\tilde{k} = 2H - \mathrm{twist}_v'$.

(2) The rotational angle of the tangent plane $\mathrm{T}_{c(t)}\,M$ is measured by the rotational speed with respect to ξ:

$$\mathrm{twist}_h' = \langle \nabla_{c'}\nu, \underbrace{c' \times \nu}_{=-Jc'}\rangle - \langle \underbrace{\nabla_{c'}\xi}_{=-HRc'}, \underbrace{c' \times \xi}_{=-Rc'}\rangle = t - H = -\tilde{k}.$$

By integrating along c we get

$$\mathrm{twist}_h = -\,\mathrm{turn}\,.$$

□

Example 4.6. We compute the torsion and the twist of a vertical geodesic c in a vertical plane in $E(\kappa, \tau)$. The geodesic c is a fibre of the Riemannian fibration. Without loss of generality $c' = \xi$, then we get

$$\begin{aligned} t &= -\nabla_{c'}\nu \cdot Jc' \\ &= \tau R\nu \cdot Jc' = -\tau. \end{aligned}$$

Namely, the torsion of a vertical geodesic is the negative of the bundle curvature.

Moreover, for the twist we get, as expected

$$\text{twist} = \int_c t + \tau l(c) = 0.$$

Hence, with respect to parallel fields, the normal does not rotate. Equivalently the normal is constant in the projection.

We shall apply Lemma 4.5 to obtain detailed information about vertical geodesics in Nil and their horizontal sister curves in $\mathbb{H}^2 \times \mathbb{R}$. This strategy is due to Laurent Mazet, for whom the author is very grateful.

For a curve $\tilde{c} \subset \mathbb{H}^2$, consider the unique horocycle foliation $\mathcal{F}_{\tilde{c}}$ given by the horocycle that is tangent to $\tilde{c}$ in $\tilde{c}(0)$ and has curvature 1 with respect to the normal n of $\tilde{c}$. Let ϑ be the angle defined by $\tilde{c}' = \cos\vartheta e_1 + \sin\vartheta e_2$, where the orthonormal frame (e_1, e_2) is given by the tangent and minus the normal of the horocycles. By the definition of the considered foliation we have $\vartheta(0) = 0$ and $n = \sin\vartheta e_1 - \cos\vartheta e_2$.



FIGURE 1. Horocycle foliation $\mathcal{F}_{\tilde{c}}$ given by $\tilde{c}$.

Lemma 4.7. *The curvature of* $\tilde{c}$ *is given by*

$$\tilde{k} = \cos\vartheta - \vartheta'.$$

Proof. A simple computation gives:

$$\begin{aligned} \nabla_{\tilde{c}'}\tilde{c}' =& \nabla_{\cos\vartheta e_1 + \sin\vartheta e_2}(\cos\vartheta e_1 + \sin\vartheta e_2) \\ =& -\vartheta' \sin\vartheta e_1 + \cos\vartheta \nabla_{(\cos\vartheta e_1 + \sin\vartheta e_2)} e_1 + \vartheta' \cos\vartheta e_2 + \sin\vartheta \nabla_{(\cos\vartheta e_1 + \sin\vartheta e_2)} e_2 \\ =& -\vartheta' \sin\vartheta e_1 + \cos^2\vartheta \underbrace{\nabla_{e_1} e_1}_{-e_2} + \cos\vartheta \sin\vartheta \underbrace{\nabla_{e_2} e_1}_{=0} \\ & + \vartheta' \cos\vartheta e_2 + \sin\vartheta\cos\vartheta \underbrace{\nabla_{e_1} e_2}_{e_1} + \sin^2\vartheta \underbrace{\nabla_{e_2} e_2}_{0} \\ =& -\vartheta' \sin\vartheta e_1 - \cos^2\vartheta e_2 + \vartheta' \cos\vartheta e_2 + \sin\vartheta\cos\vartheta e_1 \\ =& (-\vartheta' + \cos\vartheta)\sin\vartheta e_1 - (-\vartheta' + \cos\vartheta)(\cos\vartheta e_2) \\ =& (\cos\vartheta - \vartheta')n. \end{aligned}$$

□

We want to control the curve $\tilde{c}$ by means of its sister c in $M \subset E(0, 1/2) = \mathrm{Nil}$. Let $\alpha(t) = \mathrm{twist}_v$ measure the twist in $c(t)$ with respect to a basic vector field chosen such that $\alpha(0) = 0$, i.e. $\tilde{e}(c(0)) = \tilde{v}(c(0))$.

Proposition 4.8. *Let $\tilde{c} \subset \mathbb{H}^2 \times \mathbb{R}$ be the horizontal sister curve of a vertical geodesic c in $E(0, 1/2) = \mathrm{Nil}$. The angle ϑ given by the horocycle foliation $\mathcal{F}_{\tilde{c}}$ and the rotational speed α' of the minimal surface normal along c are related as follows:*

$$\vartheta' = \alpha' + \cos\vartheta - 1, \qquad \textit{and} \qquad \vartheta \leq \alpha,$$

where $\alpha(t)$ measures the angle between v and any parallel field chosen such that $\alpha(0) = 0$.

Proof. We have seen in Lemma 4.5 that $\tilde{k} = 1 - \alpha'$. Together with Lemma 4.7 we get

$$\vartheta' = \alpha' + \cos\vartheta - 1,$$

namely $\vartheta' \leq \alpha'$. In particular, along the curves c and $\tilde{c}$ we have

$$\int_{\tilde{c}} \vartheta' \leq \int_{c} \alpha' \Rightarrow \vartheta(t) \leq \alpha(t).$$

□

5. Reference surfaces

5.1. **Horizontal helicoids in** Nil_3**.** In the construction of the k-noid with genus 1 from Section 6, especially to solve one of the period problems, we use as a barrier the horizontal helicoid $H_\alpha(u, v)$ in Nil, $\left(\mathbb{R}^3, \mathrm{d}\,x_1^2 + \mathrm{d}\,x_2^2 + (\mathrm{d}\,x_3 - x_1\,\mathrm{d}\,x_2)^2\right)$ by Daniel and Hauswirth [DH09, Section 7]. For $\alpha > 0$ the coordinates of the helicoid H_α are given in terms of the solution ψ of the ordinary differential equation $\psi'^2 = \alpha^2 + \cos^2\psi$, $\psi(0) = 0$:

$$\begin{aligned} x_1 &= \frac{\sinh(\alpha v)}{\alpha(\psi'(u) - \alpha)} \cos\psi(u) \\ x_2 &= -G(u) \\ x_3 &= \frac{-\sinh(\alpha v)}{\alpha(\psi'(u) - \alpha)} \sin\psi(u), \end{aligned}$$

where G is defined by $G'(u) = 1/(\psi'(u) - \alpha)$, $G(0) = 0$. In [DH09] was shown that the function ψ is a decreasing odd bijection. There exists a unique $U := U(\alpha) > 0$ with $\psi_\alpha(U) = -\pi/2$, $\psi_\alpha(-U) = \pi/2$. To visualise the surface, we look at three curves in the helicoid:

$$\begin{aligned} H_\alpha(-U, v) &= \left(0, G(U), \frac{\sinh(\alpha v)}{\alpha(\alpha - \psi'(-U))}\right) \\ H_\alpha(0, v) &= \left(\frac{\sinh(\alpha v)}{\alpha(\psi'(0) - \alpha)}, 0, 0\right) \\ H_\alpha(U, v) &= \left(0, -G(U), \frac{\sinh(\alpha v)}{\alpha(\psi' - \alpha)}\right). \end{aligned}$$

The rulings $H_\alpha(\pm U, v)$ are vertical and define the width $a := G(-U) - G(U)$ of the helicoid. The width is well-defined for the whole helicoid, since $\psi(u + 2U) = \psi(u) - \pi$.



Figure 2. Sketch of a fundamental piece of the horizontal helicoid from Daniel and Hauswirth in Nil, $v \leq 0$.

To ensure that we can consider a helicoid H_α for a given width a, we need the following lemma:

Lemma 5.1. *For $a > 0$, there exists $\alpha > 0$ such that*

$$-2G(U(\alpha)) = a,$$

where $U(\alpha)$ is defined by $\psi_\alpha(U) = -\pi/2$. Furthermore, for $a \to 0$ we have $\alpha \to \infty$.

Proof. The idea of the proof is to show that G is a bijection on $\mathbb{R}$ as a first step. The second step is to show that U is a continuous map to $\mathbb{R}_+$.

Step 1: The function G is odd, so $G(-U) - G(U) = -2G(U)$. For $a > 0$ we show that there exists exactly one $U > 0$ such that $-2G(U) = a$: From $G' = 1/(\psi' - \alpha) < 0$ we know that G is a decreasing function on $\mathbb{R}$. If we assume that G is bounded, i.e. $G(u) \to g \in \mathbb{R}$ for $u \to \infty$, then $G'(u) \to 0$ for $u \to \infty$. But this implies $\psi'(u) - \alpha \to -\infty$ for $u \to \infty$, which is a contradiction because $\psi'^2 = \alpha^2 + \cos^2\psi \leq \alpha^2 + 1$ is bounded. Therefore, G is a decreasing bijection on $\mathbb{R}$.

Step 2: We show for $U > 0$ the existence of $\alpha > 0$ such that the solution ψ_α of

$$\psi_\alpha'^2 = \alpha^2 + \cos^2\psi_\alpha, \quad \psi_\alpha(0) = 0$$

satisfies $\psi_\alpha(U) = -\pi/2$.

By applying seperation of variables to the ODE $\psi_\alpha' = -\sqrt{\alpha^2 + \cos^2\psi}$, $\psi(0) = 0$, the solution is then given by the inverse of the elliptic integral of the first kind

$$\int_0^\psi -\frac{1}{\sqrt{\alpha^2 + \cos^2\vartheta}}\, \mathrm{d}\,\vartheta.$$

We are interested in $U(\alpha)$ given by $\psi_\alpha(U) = -\pi/2$:

$$\begin{aligned} U(\alpha) &= \int_0^{-\pi/2} -\frac{1}{\sqrt{\alpha^2 + \cos^2 \vartheta}} \, \mathrm{d}\, \vartheta \\ &= \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{\alpha^2 + \cos^2 \vartheta}} \, \mathrm{d}\, \vartheta \\ &= \frac{1}{\sqrt{\alpha^2+1}} \int_0^{\frac{\pi}{2}} \frac{1}{\sqrt{1 - \frac{1}{\alpha^2+1} \sin^2 \vartheta}} \, \mathrm{d}\, \vartheta \\ &= \frac{K\left(1/\sqrt{\alpha^2+1}\right)}{\sqrt{\alpha^2+1}}, \end{aligned}$$

where $K(k) = \int_0^{\pi/2} \frac{\mathrm{d}\,\vartheta}{\sqrt{1-k^2 \sin^2 \vartheta}} = \int_0^1 \frac{\mathrm{d}\, t}{(1-t^2)(1-k^2t^2)}$ denotes the complete elliptic integral of the first kind, defined for $k \in [0,1)$, with the special values $K(0) = \pi/2$ and $\lim_{k \to 1} K(k) = \infty$. For $\alpha \to 0$ we have $K\left(1/\sqrt{\alpha^2+1}\right) \to \infty$ and for $\alpha \to \infty$ we have $K\left(1/\sqrt{\alpha^2+1}\right) \to \pi/2$. Therefore, U is continuous because K is. Moreover, for $\alpha \to 0$ we have $U(\alpha) \to \infty$ and for $\alpha \to \infty$ we have $U(\alpha) \to 0$.

Together with the Step 1, this concludes the first part of the lemma. For the second part, notice that $a \to 0$ implies $U \to 0$, since G is a decreasing function with $G(0) = 0$. Furthermore, $U(\alpha) \to 0$ implies $\alpha \to \infty$, since K is bounded. □

We want to express the height b of the conormal η of the helicoid H_α along the vertical rulings depending on the width a and the angle φ in the horizontal plane $\mathrm{span}\{\partial x_1, \partial x_2 + x_1 \partial x_3\}$ given by

$$\cos \varphi = \langle \partial x_1, \eta \rangle.$$

The conormal along the horizontal ruling

$$H_\alpha(-U, v) = \left(0, G(U), \frac{\sinh(\alpha v)}{\alpha(\psi'(-U) - \alpha)}\right)$$

is given by

$$\frac{\partial H_\alpha}{\partial u}(-U, v) = \left(\frac{-\sinh(\alpha v)\psi'(-U)}{\alpha(\psi'(-U) - \alpha)}, -G'(-U), 0\right), \quad \nu = \frac{\partial_u H_\alpha}{\|\partial_u H_\alpha\|}.$$

The conormal along $H_\alpha(-U, v)$ is horizontal, since $x_1 = 0$. We may express φ in terms of $(\alpha, U = U(\alpha))$. By [DH09] we have

$$\psi'(-U) = G'(-U) \cos^2 \psi(-U) - \alpha = -\alpha \quad \text{and} \quad G'(-U) = \frac{1}{\psi'(-U) - \alpha} = \frac{-1}{2\alpha}.$$

Therefore

$$\eta = \frac{2\alpha^2}{\sqrt{\alpha^2(\sinh^2(\alpha v) + 1)}} \partial_u H_\alpha(-U, v) = \frac{2\alpha}{\cosh(\alpha v)} \partial_u H_\alpha(-U, v)$$

and

$$\cos\varphi = \frac{2\alpha^2\sinh(\alpha v)}{-2\alpha^2\cosh(\alpha v)} = \tanh(-\alpha v),\ v \le 0 \Leftrightarrow v = \frac{-1}{2\alpha}\ln\left(\frac{1+\cos\varphi}{1-\cos\varphi}\right).$$

Using this we get the height b of the conormal η, in terms of φ and $(\alpha, U(\alpha))$ for fixed $\alpha > 0$, as follows:

$$b = \frac{\sinh\left(\frac{-1}{2}\ln\left(\frac{1+\cos\varphi}{1-\cos\varphi}\right)\right)}{2\alpha^2} \le 0.$$

One readily sees that $\varphi \to 0$ implies $b \to -\infty$, and $b \to 0$ when $\varphi \to \pi/2$.

5.2. **Constant mean curvature k-noids in $\Sigma(\kappa)\times\mathbb{R}$.** In [GK10, Section 5.] Große-Brauckmann and Kusner constructed a one-parameter family of surfaces with constant mean curvature $H \ge 0$ in $\Sigma(\kappa)\times\mathbb{R}$, which have k ends and dihedral symmetry. They solved a Plateau problem of disc type in $E(\kappa+4H^2, H)$, its sister generates the CMC surface by reflections about horizontal and vertical planes. We use the minimal disc $M = M(a,k)$ as a barrier in our constructions.

The minimal surface M is the limit of a sequence of compact Plateau solutions $M_{(r,s)}$, which represent sections in $E(\kappa+4H^2, H)$. Each minimal disc $M_{(r,s)}$ is bounded by horizontal and vertical geodesics, see Figure 3. Let $\Gamma_{(r,s)}$ denote the boundary. The minimal surface $M_{(r,s)}$ is a section of the trivial line bundle $\pi\colon \Omega_r \subset E(\kappa + 4H^2, H) \to \Delta_r$, where $\Omega_r := \pi^{-1}(\Delta_r)$ is a mean convex domain, which is defined as the preimage of a triangle $\Delta_r \subset \Sigma(\kappa+4H^2)$. The triangle Δ_r is given by a hinge of lengths a and r, enclosing an angle π/k. The parameter a determines the length of the horizontal edge in the boundary of M, it corresponds to the necksize in the CMC sister.



FIGURE 3. The boundary of the minimal disc in $E(\kappa+4H^2, H)$.

The Jordan curves $\Gamma_{(r,s)}$ converge to the boundary Γ of the desired minimal disc M. Große-Brauckmann and Kusner proved the following theorem:

Theorem 5.2 ([GK10]). *Suppose $\kappa + 4H^2 \leq 0$. Then there exist minimal surfaces $M = M(a, k)$ with boundary Γ, having the following properties:*

- *M is a section projecting to $\Delta := \lim_{r\to\infty} \Delta_r$,*
- *M extends without branch points by the Schwarz reflection about the edges of Γ.*

6. Constant mean curvature k-noids with genus 1

We construct surfaces with MC 1/2 in $\mathbb{H}^2 \times \mathbb{R}$ with k ends and genus 1. Each surface has k vertical symmetry planes and one horizontal symmetry plane, where $k \geq 3$. The idea is to solve a Plateau problem of disc type in $\mathrm{Nil}_3(\mathbb{R})$, where the disc is bounded by geodesics. Its sister disc in $\mathbb{H}^2 \times \mathbb{R}$ generates an MC 1/2 surface by reflections about horizontal and vertical planes. The problem is to define the geodesic contour such that the sister has the desired properties.

6.1. **Boundary construction.** In $\mathbb{H}^2 \times \mathbb{R}$ the desired boundary is connected and consists of four mirror curves in three symmetry planes; the two vertical symmetry planes form an angle π/k, see Figure 4.



Figure 4. The desired boundary of the 1/2-surface in $\mathbb{H}^2 \times \mathbb{R}$ and its minimal sister surface in Nil.

By [GK10] the sister surface is bounded by a geodesic contour $\Gamma := \Gamma(a, b, \varphi)$: The horizontal mirror curves correspond to vertical geodesics and the vertical mirror curves correspond to horizontal geodesics, their projections enclose an angle $\varphi \in (0, \pi)$.

The length $a > 0$ of the finite horizontal geodesic c_1 determines the asymptotic parameter of the ends of the k-noid, called the necksize. The length $b > 0$ of the vertical finite edge c_2 defines the size of the hole of the k-noid: For $b \to 0$ the k-noid is close to the non-degenerate k-noid of [GK10]. The angle φ measures the curvature of $\tilde{c}_2$. The parameters will be determined by solving the period problems.

To construct a minimal surface that is bounded by Γ, we truncate the infinite contour Γ and get Jordan curves Γ_n, $n > 0$. We consider a geodesic triangle Δ_n in the base manifold $\mathbb{R}^2$ of the Riemannian fibration of $\mathrm{Nil}_3(\mathbb{R})$: Two edges of lengths a and n form an angle φ and intersect in a point $\hat{p}_1$. We lift $\hat{p}_1$ and the corresponding edge of length n horizontally and label the vertices with p_1 and p_6. Then we add a vertical arc of length b at p_1 in fibre-direction ξ and label its endpoint with p_2. We lift the edge of length a of the base triangle in $\mathbb{R}^2$ horizontally, such that it ends in p_2; the other vertex is labelled by p_3. We add another vertical edge in fibre-direction ξ: it starts in p_3, has length n^2 and its end vertex is called p_4. By lifting the remaining edge of the base triangle horizontally and inserting another vertical edge with endpoints p_5 and p_6 we complete the special Jordan curve Γ_n.

In general the vertical distances do not sum up: $d(p_1, p_2) + d(p_3, p_4) \neq d(p_5, p_6)$, but we claim p_6p_5 is in fibre direction. We consider an arc length parametrization γ of $\partial\Delta_n \subset \mathbb{R}^2$ which runs counter-clockwise. By Lemma 2.4 the vertical distance of its horizontal lift is $\mathrm{area}(\Delta_n)$. By construction we get

$$d(p_6, p_5) = b + n^2 - \mathrm{area}(\Delta_n).$$

Since $\mathrm{area}(\Delta_n)$ grows linearly, there exists $N \in \mathbb{N}$ such that for all $n \geq N$: $d(p_6, p_5) > 0$ and $d(p_6, p_5) \to \infty$ for $n \to \infty$.

The polygon Γ_n has six right angles; its projection Δ_n is convex for every n and has one fixed angle $\varphi \leq \pi$ independent of n. We define a mean convex set $\Omega_n := \pi^{-1}(\Delta_n) \subset \mathrm{Nil}_3(\mathbb{R})$.

For $n \to \infty$ we have $\Gamma_n \to \Gamma$. To control the Plateau solution for Γ, we solve the Plateau problem for Γ_n and consider their limit.

Lemma 6.1. *The special Jordan curve $\Gamma_n \subset \mathrm{Nil}_3(\mathbb{R})$ bounds a unique Plateau solution $M_n \subset \Gamma_n$. It is a section over Δ_n and extends without branch points by Schwarz reflection about the edges of Γ_n.*

Proof. Since Δ_n is convex and $\partial\Delta_n$ consists of geodesics, its preimage $\Omega_n = \pi^{-1}(\Delta_n)$ is a mean convex set and by construction we have $\Gamma_n \subset \partial\Omega_n$. Therefore, the Plateau problem is solvable by [HS88]. Moreover, by Section 3 the solution M_n is a unique section of the trivial line bundle $\pi\colon \Omega_n \to \Delta_n$ that extends as an immersion across Γ_n. ☐

6.2. **Plateau solutions.** The Plateau solutions M_n are contained in a nested sequence of mean convex sets, hence we are able to solve the Plateau problem for Γ:

Theorem 6.2. *There exists a unique minimal surface $M(a, b, \varphi) \subset \mathrm{Nil}_3(\mathbb{R})$, which is a section over $\Delta := \pi(\lim \Gamma_n)$ and extends without branch points by Schwarz reflection about its edges for all $a, b > 0$ and $\varphi \in (0, \pi)$.*

Proof. Two copies of the minimal k-noid from [GK10] bound the sequence M_n from below and above if they coincide along the vertical axis and are obtained by a vertical translation. Therefore there exists a uniform curvature estimate and a limit by [Gro93]. Thus, there exists a minimal surface $M(a, b, \varphi)$ of disc-type for all $a, b > 0$ and $\varphi \in (0, \pi)$, which is regular in the interior and $\partial M = \Gamma$. Since Γ consists of geodesics and has bounded geometry, Schwarz reflection applies and M is smooth at the boundary.

By Proposition 3.3 the limit is a section, since $\lim \Gamma_n$ is not contained in a vertical plane. ☐

6.3. **Period problems.** To construct an MC 1/2 surface with genus 1 and certain symmetries we have to solve two period problems. One period is given by the vertical distance of the two horizontal mirror curves. The second period is angular and determined by the geometry of the finite horizontal curve.

To solve the first period problem for the MC 1/2 surface $\tilde{M} \subset \mathbb{H}^2 \times \mathbb{R}$, i.e. to construct a minimal surface $M \subset \mathrm{Nil}_3(\mathbb{R})$ such that the two horizontal components of its sister surface lie in the same horizontal plane, we consider the mirror curve in the vertical plane with finite length a in $\partial\tilde{M}$, and call it $\tilde{c}_1$. The period is given by $p = \int \langle \tilde{c}_1', \tilde{\xi} \rangle_{\mathbb{H}^2\times\mathbb{R}}$, where $\tilde{\xi}$ is the vertical vector field of $\mathbb{H}^2 \times \mathbb{R}$. Since we have a first order description for the vertical parts of vector fields, we consider the vector field $\tilde{T}$ on $\tilde{M}$ given by $\tilde{\xi} - \langle \tilde{\xi}, \tilde{\nu} \rangle \tilde{\nu}$, where $\tilde{\nu}$ denotes the normal of $\tilde{M}$. It is the tangential projection of the vertical vector field and rotates by $\pi/2$ in the tangent plane under conjugation:

$$\mathrm{d} f^{-1}(T) = J \,\mathrm{d} \tilde{f}^{-1}(\tilde{T}),$$

where f and $\tilde{f}$ denote the corresponding parametrisations of M and $\tilde{M}$. Therefore, we have an analogous formulation of the period on M:

$$\begin{aligned} p(\tilde{M}) &= \int_{\tilde{c}_1} \langle \tilde{c}_1', \tilde{\xi} \rangle_{\mathbb{H}^2\times\mathbb{R}} = \int \langle \mathrm{d}\tilde{f}(\gamma'), \tilde{T} \rangle_{\mathbb{H}^2\times\mathbb{R}} \\ &= \int \langle \mathrm{d}\tilde{f}(\gamma'), \mathrm{d}\tilde{f}(J^{-1}\,\mathrm{d} f^{-1}(T)) \rangle_{\mathbb{H}^2\times\mathbb{R}} = \int \langle J\gamma', \mathrm{d} f^{-1}(T) \rangle \\ &= \int_{c_1} \langle \eta, \xi \rangle_{\mathrm{Nil}_3(\mathbb{R})} = p(M). \end{aligned}$$

As seen above, we have a Plateau solution $M(a, b, \varphi)$ for all $a, b > 0$ and $\varphi \in (0, \pi)$. The first period problem is solvable for each $a > 0$ and $\varphi \in (0, \pi/2)$:

Proposition 6.3. *For each $a > 0$ and $\varphi \in (0, \pi/2)$ there exists $b(a, \varphi) > 0$ such that $p(M(a, b(a, \varphi), \varphi)) = 0$.*

Proof. Let $a > 0$ and $\varphi \in (0, \pi/2)$ be fixed. By Theorem 6.2 we have a unique Plateau solution $M_b := M(a, b, \varphi)$ for each $b > 0$. We claim the function $p(b) := p(M_b)$ is continuous in b. To see this, we take two converging sequences (b_l) and (b_k) with the same limit b_0. If $\lim p(b_l) \neq \lim p(b_k)$ this would contradict the uniqueness of the minimal section, since the corresponding minimal surfaces also converge: There is a uniform curvature estimate for each sequence of minimal surfaces M_{b_l} by [GK10]. Moreover since the surfaces represent sections, each sequence satisfies a uniform area estimate. Writing the surfaces as graphs in normal coordinates over their tangent planes, we see that the sequence contains a converging subsequence with limit $M = M_{b_0}$.

We will now show that there exists $b_t \in \mathbb{R}$ such that $p(b) < 0$, for all $b > b_t$, and $\lim_{b\to 0} p(b) > 0$. By the intermediate value theorem this proves the lemma.

- To define b_t, we consider the horizontal helicoid M_H constructed by Daniel and Hauswirth [DH09] from Section 5.1, whose horizontal axis coincides with c_1. By Lemma 5.1 there exists a helicoid M_H for each pitch $a > 0$ such that the incident vertical arcs of Γ are contained in its rulings. Since $\varphi < \pi/2$ the helicoid M_H and M_b are both minimal sections over a bounded convex domain $\pi(M_H) \cap \pi(M_b)$, whose boundary consists of three geodesic arcs.

We claim there exists $b_t > 0$, such that the surfaces intersect in Γ only. In Section 5.1 we showed that the conormal η_H of the helicoid along the vertical ruling is horizontal and its opening angle depends continuously on the height given by

$$h = \frac{\sinh\left(\frac{-1}{2}\ln\left(\frac{1+\cos\varphi}{1-\cos\varphi}\right)\right)}{2\alpha^2},$$

where α depends on the pitch a, see Lemma 5.1. We consider the vertical plane V, given by its tangent plane, that is spanned by the conormal η_H and ξ in height h. Since each conormal is horizontal and turns monotonically, the intersection $V \cap M_H$ meets the vertical ruling in exactly one point given by the height h. Moreover M_H is a section in the interior and therefore $V \cap M_H$ is bounded from below. Hence for $a > 0$ and $\varphi \in (0, \pi/2)$ exists $b_t \geq |h|$ such that the helicoid is a barrier lying above M_b for every $b > b_t$.

Consequently we can estimate the vertical component of the conormal η of the Plateau solution M_b by the helicoid conormal η_H along the interior of the curve c_1:

$$\langle \eta, \xi \rangle < \langle \eta_H, \xi \rangle \quad \text{and hence} \quad p(M_b) = \int \langle \eta, \xi \rangle < \int \langle \eta_H, \xi \rangle = 0.$$

- Otherwise for $b \to 0$ we consider a minimal surface N in $\mathrm{Nil}_3(\mathbb{R})$ defined in [GK10, Section 5.], see Section 5.2 with the parameters a and φ. It has a positive period p_N, since it is bounded from below by a horizontal umbrella.

  For $b \to 0$ we have $M_b \to N$ away from the singularity. Furthermore, the sequence of conormals η_b converges uniformly to η_N on compact sets $K \subset c_1$. Therefore, the period $p(b)|_K$ converges uniformly to $p_N|_K > 0$ for each compact $K \subset c_1$ and $b \to 0$. Hence, on c we have $p(b) > 0$ for $b \to 0$.

□

Remark 6.4. For $\varphi = \pi/2$ the proof does not work since the helicoid M_H is not a barrier for $b < \infty$. Therefore, we cannot construct a genus 1 catenoid in $\mathbb{H}^2 \times \mathbb{R}$ with this method.

Lemma 6.5. *For each $a > 0$ and $\varphi \in (0, \pi/2)$ the map $b\colon \mathbb{R}_+ \times (0, \pi/2) \to \mathbb{R}_+$ defined by Proposition 6.3 is a continuous function.*

Proof. We assume that b is discontinuous in (a, φ), i.e. there exist two sequences (a_l, φ_l) and $(\bar{a}_l, \bar{\varphi}_l)$ with limit (a_0, φ_0) but w.l.o.g. $b_0 = \lim b(a_l, \varphi_l) > \lim b(\bar{a}_l, \bar{\varphi}_l) = \bar{b}_0$. There is a uniform curvature estimate for each sequence of minimal surfaces $M(a_l, b(a_l, \varphi_l), \varphi_l)$ and $M(\bar{a}_l, b(\bar{a}_l, \bar{\varphi}_l), \bar{\varphi}_l)$ by [GK10]. Moreover since the surfaces represent sections, each sequence satisfies a uniform area estimate. Writing the surfaces as graphs in normal coordinates over their tangent planes, we see that each one contains a converging subsequence with limit $M = M(a_0, b_0, \varphi_0)$ and $\bar{M} = M(a_0, \bar{b}_0, \varphi_0)$ respectively. Both minimal surfaces M and $\bar{M}$ have zero period. But for $b_0 > \bar{b}_0$ the minimal surface M bounds a mean convex domain from above with $\partial\bar{M}$ in its boundary. Therefore, M is an upper barrier for $\bar{M}$ with $\langle \eta, \xi \rangle > \langle \bar{\eta}, \xi \rangle$, contradicting the fact that both minimal surfaces have zero period. □

We solved the first period problem in b depending on (a, φ). Namely, for each angle $\varphi \in (0, \pi/2)$ and horizontal geodesic of length a, there exists an $b > 0$ as length of the vertical geodesic, such that the two horizontal mirror curves in the sister surface lie in the same mirror plane.

The second period problem relies on the horizontal mirror curve $\tilde{c}_2$ of length b and there are two difficulties, we have to solve: First of all, we have to make sure that the two vertical mirror planes, which are perpendicular to $\tilde{c}_2$ intersect. And secondly their angle of intersection has to be π/k. For the answers we need to restate the solution of the first period problem: For an angle φ and a vertical geodesic c_2 of length b there exists a horizontal geodesic of length a, such that the period is zero:

Proposition 6.6. *For each $b > 0$ and $\varphi \in (0, \pi/2)$ there exists $a(b, \varphi) > 0$, such that the horizontal period of the minimal surface $M(a(b, \varphi), b, \varphi)$ is zero.*

Proof. By Lemma 6.5 the map $b\colon \mathbb{R}_+ \times (0, \pi/2) \to \mathbb{R}_+$ continuous. For each φ we claim $b(a, \varphi) \to 0$ for $a \to 0$. Indeed, by the proof of Proposition 6.3 there exists $-h \in \mathbb{R}$ as an upper bound of $b(a, \varphi)$ given by a and φ:

$$-h = \frac{\sinh\left(\frac{1}{2}\ln\left(\frac{1+\cos\varphi}{1-\cos\varphi}\right)\right)}{2\alpha(a)^2} \geq b(a, \varphi), \quad \text{since otherwise} \quad p > 0.$$

By Lemma 5.1 we know that $\alpha(a) \to \infty$ for $a \to 0$. Therefore, the height h converges to zero as well as $b(a, \varphi)$ converges to zero.

Remains to show that the map b is unbounded. For each $\varphi > 0$ assume the contrary: The function $b_\varphi(a) := b(a, \varphi) \leq \hat{b}$ is bounded. Since b is a continuous function, we have $p(M_{\tilde{b}}) \neq 0$ for $\tilde{b} > \hat{b}$. The minimal surface $M(a, b(a, \varphi), \varphi)$ is a barrier from above for $M(a, \tilde{b}, \varphi)$, therefore $p(M_{\tilde{b}}) < 0$. The continuity of p in b implies $p(M_{\tilde{b}}) < 0$ for $\tilde{b} > \hat{b}$ and for all $a > 0$.

Consider the map $p_1 \colon a \mapsto p(M(a, \tilde{b}, \varphi))$. We claim the function p_1 is continuous. To see this, we take two converging sequences (a_l) and (a_k) with the same limit a_0. Since the corresponding minimal surfaces also converge, this implies $\lim p(a_l) = \lim p(a_k)$. Moreover, for a large enough the minimal surface has a positive period, therefore there exists $\hat{a} > 0$ with $p_1(\hat{a}) = 0$, which is a contradiction.

Hence, $b_\varphi(a)$ is unbounded with $b_\varphi(a) \to 0$ for $a \to 0$. So we conclude: For all $b > 0$ there exists $a(b, \varphi) > 0$ (not necessarily unique) such that $M(a(b, \varphi), b, \varphi)$ has zero period. □

Remark 6.7. For conjugate minimal surfaces in $\mathbb{R}^3$ we have a unique $a(b, \varphi)$, since $a \mapsto b(a, \varphi)$ is injective because of scaling.

Before we solve the second period problem, we want to analyse the finite horizontal mirror curve $\tilde{c}_2$ in $\mathbb{H}^2 \times \mathbb{R}$ parametrized by arc length. We choose the downward-pointing surface normal ν and (c_2', η, ν) positive oriented. Where c_2 is the sister curve, then $\langle c_2', \xi \rangle = 1$. We consider the twist $\varphi(t)$ of c_2. The curvature of $\tilde{c}_2$ is $\tilde{k} = 1 - \varphi'(t) < 1$, since $\varphi' > 0$ by graph property of the minimal surface. Proposition 4.8 implies the embeddedness, since $\vartheta \leq \varphi < \pi/2$; moreover, $\vartheta \geq 0$ since $\vartheta < 0$ implies $\tilde{k} > 1$. With γ_0 and γ_b we denote the unique geodesics given by $\gamma_i(0) = \tilde{c}_2(i)$ and $\gamma_i'(0) = -\tilde{\nu}(i)$, $i = 0, b$, see Figure 5.

As said before, in a first step we have to make sure, that the geodesics intersect:



FIGURE 5. Sketch of the defined geodesics γ_i, $i = 0, b$.

Proposition 6.8. *For each* $\varphi \in (0, \pi/2)$ *exists* $b_0 := b_0(\varphi) > 0$ *satisfying*

$$b_0 e^{b_0} = \frac{1 - \cos(\varphi - b_0)}{\sin(\varphi - b_0)}, \quad b_0(\varphi) < \varphi$$

such that the vertical mirror planes of $\tilde{M}(a(b, \varphi), b, \varphi)$ *intersect for all* $b \in (0, b_0(\varphi))$ *and define an intersection-angle* $\alpha > 0$.

Proof. As in Proposition 4.8 we consider the foliation by horocycles given by $\tilde{v}(0)$ and the related angle $\vartheta \leq \varphi < \pi/2$. For the calculation we consider the upper halfplane and orient $\tilde{c}_2$ such that $\tilde{c}_2(0) = (0, 1)$ and $\tilde{v}(0) = (0, 1)$, then γ_0 is contained in the y-axis. We want to parametrize the unique geodesic $\gamma_b \subset \mathbb{H}^2$, which starts in the endpoint of $\tilde{c}_2$ ($\tilde{c}_2(b) = (c_x, c_y)$) and its tangent is parallel to $(\sin(\vartheta), -\cos(\vartheta))$. For $\vartheta \neq k\pi, k \in \mathbb{Z}$, γ_b is an Euclidean halfcircle with radius r and its midpoint on the x-axis. We solve the linear equations

$$\gamma_b(\pi - \vartheta) = (c_x, c_y) = (x + r\cos(\pi - \vartheta), r\sin(\pi - \vartheta)).$$

The geodesic is in Euclidean coordinates parametrised by

$$\gamma_b(t) = \left(c_x + c_y\left(\frac{\cos(\pi - t) - \cos\vartheta}{\sin\vartheta}\right), c_y\frac{\sin(\pi - t)}{\sin\vartheta}\right).$$

The geodesics intersect if the x-coordinate of $\gamma_b(\pi)$ is positive:

$$c_x + c_y\frac{1 - \cos\vartheta}{\sin\vartheta} > 0. \tag{6.1}$$

Since $d_{\mathbb{H}^2}(\tilde{c}_2(0), \tilde{c}_2(b)) \leq b$ we have $c_x \geq -b$ and $c_y > e^{-b}$. So we conclude Equation (6.1) is true if

$$\frac{1 - \cos\vartheta}{\sin\vartheta} > be^b.$$

The angles ϑ and φ are related by $\vartheta' = \varphi' + \cos\vartheta - 1$ (see Proposition 4.8) and $\cos\vartheta \geq 0$ implies $\int(\cos\vartheta - 1) \geq -b$. Therefore we know that $\vartheta \geq \varphi - b$. Furthermore, the function $\vartheta \mapsto (1 - \cos\vartheta)/\sin\vartheta$ increases monotonically, so we conclude, the geodesics intersect if

$$be^b < \frac{1 - \cos(\varphi - b)}{\sin(\varphi - b)}.$$

This is equivalent to

$$f_\varphi(b) := \frac{1 - \cos(\varphi - b)}{\sin(\varphi - b)} - be^b > 0. \tag{6.2}$$

Its differential

$$f'_\varphi(b) = \frac{\cos(\varphi - b) - 1}{\sin^2(\varphi - b)} - e^b(1 + b)$$

is less than zero for all $b < \varphi$. Hence, f_φ is decreasing. Let us consider the limits at the boundaries:

$$\lim_{b\to 0} f_\varphi(b) = \frac{1 - \cos\varphi}{\sin\varphi} > 0, \qquad \text{for } \varphi \in (0, \pi/2)$$
$$\lim_{b\to\varphi} f_\varphi(b) = -\varphi e^{\varphi} < 0.$$

We conclude, there exists exactly one $b_0(\varphi) \in (0, \varphi)$ with $f_\varphi(b_0(\varphi)) = 0$. Moreover, for all $b < b_0(\varphi)$ we have $f_\varphi(b) > 0$, therefore the geodesics intersect for all $b < b_0(\varphi)$. □

Remark 6.9. For $b = \varphi$ the total curvature of $\tilde{c}_2$ is $b - \varphi = 0$ and therefore the two geodesics γ_0 and γ_b do not intersect in any $p \in \mathbb{H}^2$, but in $\partial\mathbb{H}^2$. By Proposition 6.6 there exists $a > 0$ to solve the first period problem, hence after reflection the contruction causes a complete singly periodic MC-1/2 surface $\tilde{M}(a(\varphi, \varphi), \varphi, \varphi)$ in $\mathbb{H}^2 \times \mathbb{R}$ with infintely many ends.

Now we are able to solve the second period problem that is given by the angle α. We want the surface to close after $2k$ reflections about vertical mirror planes, so α has to be π/k.

Proposition 6.10. *For each $k \geq 3$ there exists $\epsilon = \epsilon(k) > 0$, such that $\varphi = \pi/k + \epsilon < \pi/2$ and there exists $0 < b < b_0(\varphi)$, such that the surface $\tilde{M}(a(b, \varphi), b, \varphi)$ has angular period $\alpha = \pi/k$ and $p(\tilde{M}(a(b, \varphi), b, \varphi)) = 0$.*

Proof. For $b < b_0(\varphi)$ the geodesics γ_0 and γ_b intersect by Proposition 6.8. Hence, we can apply the Gauß-Bonnet Theorem to the compact disc $V \subset \mathbb{H}^2$ defined by $\partial V = \tilde{c}_2 \cup \gamma_0 \cup \gamma_b$:

$$\int_V K + \int_{\partial V} k_g + \sum \alpha_i = 2\pi\chi(V),$$

where α_i, $i = 1, 2, 3$ are the exterior angles with $\alpha_i = \pi/2$ for $i = 1, 2$ and $\alpha_3 = \pi - \alpha$. Furthermore, k_g is the geodesic curvature with respect to the inner normal of ∂V and since $\tilde{c}_2$ is a mirror curve $k_g = -\tilde{k}$ with respect to the surface normal. Using this we get

$$\int_V K - \int_{\tilde{c}_2} \tilde{k} + 2\pi - \alpha = 2\pi. \tag{6.3}$$

From Lemma 4.5 we know $\tilde{k} = 1 - \varphi'$. Integrating this shows Equation (6.3) is equivalent to $\varphi - b - \operatorname{area}(V) = \alpha$.

We claim, that the lengths $l(\gamma_i)$ are bounded from above for all $b \leq b_0(\varphi)$. Recall from the proof of Proposition 6.8 that $\varphi \geq \vartheta \geq \varphi - b$. In particular, we have a lower bound for ϑ, the angle that is given by the tangent of $\tilde{c}_2$ and the horocycle fibration. Assume that the lengths $l(\gamma_i)(b) \to \infty$ for $b \to 0$, this implies $\vartheta \to 0$, a contradiction. If the lengths are bounded, the area tends to zero for $b \to 0$.

The angle α depends continuously on $b < b_0(\varphi)$:

$$\alpha(b) = \varphi - b - \operatorname{area}(V(b)) \tag{6.4}$$

and decreases. The idea is to show that for any $k \geq 3$ exists $\varphi_k \in (0, \pi/2)$ such that

$$\lim_{b \to 0} \alpha(b) > \pi/k \quad \text{and} \quad \lim_{b \to b_0(\varphi_k)} \alpha(b) < \pi/k.$$

On the one hand $\lim_{b \to 0} \alpha(b) = \varphi_k$, hence we have to choose $\varphi_k > \pi/k$. Therefore, for any $\epsilon_k > 0$, $\varphi_k := \pi/k + \epsilon_k$ satisfies the first condition.

On the other hand we have

$$\lim_{b \to b_0(\varphi_k)} \alpha(b) = \varphi_k - b_0(\varphi_k) - \operatorname{area}(V(b_0(\varphi_k))),$$

hence we have to choose $\epsilon_k > 0$ such that

$$\varphi_k < \frac{\pi}{k} + b_0(\varphi_k) + \operatorname{area}(b_0(\varphi_k)) \quad \Leftrightarrow \quad \epsilon_k < b_0(\varphi_k) + \operatorname{area}(b_0(\varphi_k)).$$

We claim the function $b_0(\varphi)$ increases monotonically. Recall Equation (6.2): b_0 was defined implicitly by $f_\varphi(b) = \frac{1-\cos(\varphi-b)}{\sin(\varphi-b)} - be^b = 0$. By chain rule we have

$$b_0'(\varphi) = e^{b_0(\varphi)}(1 + b_0(\varphi)) + \frac{1 - \cos(\varphi - b_0(\varphi))}{\sin^2(\varphi - b_0(\varphi))} > 0.$$

Therefore with $\epsilon_k = b_0(\pi/k)$ we get:

$$\lim_{b \to b_0(\varphi_k)} \alpha(b) = \pi/k + \underbrace{b_0(\pi/k) - b_0(\varphi_k)}_{<0} - \operatorname{area}(V(b_0(\varphi_k))) < \pi/k.$$

Remains to show that $\varphi_k < \pi/2$ for all $k \geq 3$. Since b_0 increases this is true if $\pi/3 + b_0(\pi/3) < \pi/2$, i.e. $b_0(\pi/3) < \pi/6$. But this follows directly from the fact that $f_{\pi/3}(\pi/6) < 0$.

By the intermediate value theorem follows, there exists $b^* \in (0, b_0(\pi/k + \epsilon_k))$ such that $\alpha(b^*) = \pi/k$. By Proposition 6.6 there exists $a > 0$ to solve the first period problem. □

The proposition proves the existence of one surface $M(a_k, b_k, \varphi_k)$ for any $k \geq 3$. It is natural to ask if there is a family of CMC surfaces with the desired symmetries and k ends. The answer is yes:

Proposition 6.11. *For each $k \geq 3$ there exists an interval $U_k \subset (0, \pi/2)$, such that for all $\varphi \in U_k$ there exists $b(\varphi) > 0$, such that the surface $\tilde{M}(a(b(\varphi), \varphi), b(\varphi), \varphi)$ has angular period $\alpha = \pi/k$ and $p(\tilde{M}(a(b, \varphi), b, \varphi)) = 0$.*

Proof. In Proposition 6.10 the existence of a triple $(a(b(\varphi_k), \varphi_k), b(\varphi_k), \varphi_k)$ was proven such that the surface $\tilde{M}(a(b(\varphi_k), \varphi_k), b(\varphi_k), \varphi_k)$ has the desired properties. By Equation (6.4) we know, the pair $(b(\varphi_k), \varphi_k)$ is a zero of the following continuously differentiable function

$$G(b, \varphi) = \alpha(b, \varphi) - \pi/k = \varphi - b - \operatorname{area}(V(b, \varphi)) - \pi/k.$$

The angle $\alpha(b, \varphi)$ is given by the two geodesics γ_0 and γ_b. Recall from the proof of Proposition 6.10 that $\partial_b \alpha(b, \varphi) < 0$. By the implicit value theorem there exist an open neighborhood U_0 of φ_k, an open neighborhood V of $b(\varphi_k)$, and a unique continuously differentiable function $g \colon U_0 \to V$ with $g(\varphi_k) = b(\varphi_k)$ such that $G(g(\varphi), \varphi) = 0$ for all $\varphi \in U_0$.

To define U_k notice that $g(\varphi_k) < b_0(\varphi_k)$, therefore the subset $U_k := \{\varphi \in U_0 \colon g(\varphi) < b_0(\varphi)\} \cap (0, \pi/2)$ is not empty. Hence, for all $\varphi \in U_k$ there exists $b = g(\varphi) < b_0(\varphi)$ and therefore by Proposition 6.6 there exists $a > 0$ to solve the first period problem. □

Remark 6.12. We want to analyse the limiting cases:

- $\varphi \to \inf U_k$: We know that $\inf U_k \geq \pi/k$. From $\varphi \to \pi/k$ follows $b + \text{area}(V(b,\varphi)) \to 0$, which implies $b \to 0$. Assuming the solution of the first period problem $a(b,\varphi) > 0$ for $b \to 0$ leads to the k-noid from [GK10] which has a positive period. Therefore if $\inf U_k = \pi/k$ then the sequence of k-noids converges to an union of k horocylinders away from the singularity for $\varphi \to \pi/k$.
- $\varphi \to \sup U_k \leq \pi/2$: Since $\partial_\varphi \,\text{area}(V(b,\varphi)) \leq 0$, $\varphi \to \sup U_k$ implies that b increases. For the CMC surfaces this means that the length of the finite horizontal symmetry curve grows.

6.4. **Main Theorem.** After solving the two period problems we can now prove the existence of the MC 1/2 surface with genus 1:

Theorem 6.13. *For $k \geq 3$, there exists a family of surfaces M with constant mean curvature $1/2$ in $\mathbb{H}^2 \times \mathbb{R}$ such that:*
• M is a proper immersion of a torus minus k points,
• M is Alexandrov embedded.
• M has k vertical mirror planes enclosing an π/k-angle,
• M has one horizontal mirror plane.

Proof. By [Dan07] the fundamental piece $M(a, b_0)$ has a sister surface $\tilde{M}$ with constant mean curvature $1/2$ in $\mathbb{H}^2 \times \mathbb{R}$, which is a graph. By construction and from the solution of the period problems, $\tilde{M}$ has one horizontal and two vertical mirror planes; the two vertical mirror planes enclose an angle π/k. It consists of four mirror curves: two horizontal (one bounded and one unbounded) and two vertical (one bounded and one unbounded). After Schwarz reflection about the vertical infinite mirror curve followed by reflection about the horizontal mirror plane we have the fundamental domain of one end: It is built up of four fundamental domains $\tilde{M}$. We use the Euler characteristic

$$\chi = V - E + F = 2 - 2g$$

to determine the genus g of the complete MC 1/2 surface M with k ends, which is generated by Schwarz reflection. We have $\chi = 4k - 8k + 4k = 0$ and therefore $g = 1$.

We claim that M is Alexandrov-embedded if we choose as before the downward-pointing normal ν. We show the fundamental piece $\widetilde{M}$ is embedded and stays in the subset of $\mathbb{H}^2 \times \mathbb{R}$ that is bounded by mirror planes. The mirror planes are given by the symmetry curves with infinte length. We discuss the embeddedness of each boundary arc, since $\widetilde{M}$ is transverse to the fibres, the fundamental domain is embedded. Therefore, the complete surface M is Alexandrov-embedded.

Let us recall the notation, c_1 denotes the finite horizontal geodesic in $\partial M(a, b_0)$ and $\tilde{c}_1$ its sister curve in $\partial\widetilde{M}$. The sister curve $\tilde{c}_1$ is a mirror curve in a vertical plane. The curve is graph and therefore embedded. The same holds for the other horizontal curve and its sister curve in a vertical plane, call it $\tilde{c}_3$.

Along the horizontal geodesic c_3 we have $\langle \eta, \xi \rangle > 0$ if we choose $(c_3'\eta, \nu)$ positive oriented. By the first order description of the sister surfaces we know that the projection of the vertical vector field ξ on the tangent plane rotates by $\pi/2$ under conjugation. Therefore, we have $\langle \tilde{c}_3', \tilde{\xi} \rangle < 0$. By construction, we know that $\langle \eta, \xi \rangle \to 1$ in the end along the sister curve c_3. Hence, in the sister surface for



the corresponding mirror curve we have $\langle \tilde{c}_3', \tilde{\xi} \rangle \to -1$. In other words, the mirror curve comes from ∞ in the end.

With the arguments of Hauswirth, Rosenberg and Spruck in the proof of Theorem 1.2 in [HRS08] we know that each divergent sequence (p_n) in M with $\langle \tilde{\nu}(p_n), \tilde{\xi} \rangle \to 0$ has a limit in the boundary in the projection: $\pi(p_n) \to \partial\mathbb{H}^2$. The idea of the proof is to show that if the sequence has a horizontal normal in the limit and a limiting point in the projection, then the whole surface is converging to a horocylinder and stays on one side. Hence, by the half-space theorem it is a horocylinder which is a contradiction.

As in the setup let c_2 denote the finite vertical geodesic, recall that its sister curve is embedded. Let c_0 denote the remaining vertical geodesic in the boundary $\partial M(a, b_0)$ parametrized in ξ-direction. Let α denote the twist of the horizontal normal ν along c_0. We have $\alpha' > 0$, i.e. the normal rotates monotonically, because M_∞ is a section. We consider the horocycle fibration of $\mathbb{H}^2$ given by $\tilde{\nu}(0)$ as in Proposition 4.8. By the proposition the angle ϑ the sister curve encloses with the fibration in $\mathbb{H}^2$ is smaller than $\alpha = \pi - \varphi < \pi$. Hence, $\tilde{c}_0$ is embedded. In summary this proves the first part, $\widetilde{M}$ is embedded.

In the second step we show that $\widetilde{M}$ is bounded by symmetry planes. First, we have to check if the surface stays in a horizontal halfspace of $\mathbb{H}^2 \times \mathbb{R}$. We define the horizontal mirror plane of M to be $\mathbb{H}^2 \times \{0\}$. By the maximum principle $\widetilde{M}$ has no (global) minimum below $\mathbb{H}^2 \times \{0\}$. Assume there is a curve $\tilde{c}$ in $\widetilde{M}$ with $\tilde{c}(0) \in \partial\widetilde{M}$, whose third component goes to minus infinity. Since the Gauß map is open, we have $\langle \tilde{c}'(t), \tilde{\xi} \rangle < 0$ for $t > 0$. But $\langle \tilde{c}', \tilde{\xi} \rangle \leq 0$ implies $\tilde{c}(0) \in \tilde{c}_1 \cup \tilde{c}_3$, since $\langle \tilde{\eta}, \tilde{\xi} \rangle = 1$ along $\tilde{c}_0 \cup \tilde{c}_2$. Consider the sister curve $c \subset M(a, b_0)$, by the first order description we have $\langle \eta, \xi \rangle \geq 0$ along c and $\langle \eta, \xi \rangle \to 1$ in the end. Moreover, since $\tilde{\nu}$ rotates within a vertical plane along $\tilde{c}_1$ and $\tilde{c}_3$ we have $0 = \langle \tilde{c}'(0), \tilde{\xi} \rangle = -\langle \eta(0), \xi \rangle$. By the maximum principle the tangent plane along c_3 is nowhere horizontal and therefore $c(0) \in c_1$. Along c_1 there exists a unique $p \in c_1$ where the tangent plane is horizontal, since the normal rotates monotone. The horizontal umbrella U_p in p intersects the minimal surface $M(a, b_0)$ in $2k$ analytic curves. In fact $k = 2$, by the monotone rotation of the normal; this implies $U_p \cap M(a, b_0) = c \cup c_1$. And $M(a, b_0)$ is below the umbrella to the side of the conormal along c, which contradicts the fact that $\langle \eta, \xi \rangle \geq 0$ along c. Hence $\widetilde{M} \subset \mathbb{H}^2 \times \mathbb{R}_0^+$.

To finish the proof we have to show that the surface lies on one side of the vertical symmetry plane: We have seen that the projection of the vertical symmetry curve $\tilde{c}_3$ is converging to some point in the boundary $\partial\mathbb{H}^2$. Along $\tilde{c}_0$ the surface is at least locally to the side of the normal, since $\tilde{k} = 1 - \alpha' < 1$ by the graph property of the minimal surface along c_0. Therefore, since it is also a graph, it is defined on a domain $\Omega \subset \mathbb{H}^2$ given by $\pi(\tilde{c}_3) \cup \tilde{c}_0$. $\square$

6.5. **Conclusion and outlook.** We constructed an MC 1/2 surface in $\mathbb{H}^2 \times \mathbb{R}$ with k ends, genus 1 and k-fold dihedral symmetry, $k \geq 3$, which is Alexandrov embedded. We had to solve two period problems in the construction. The first period guarantees that the surface has exactly one horizontal symmetry. For the second period we had to control a horizontal mirror curve to get the dihedral symmetry. In the case of $H \neq 0$ the total curvature of a horizontal mirror curve depends not only on the twist of the normal, but also on its length. An interesting problem is to construct non-embedded examples that correspond to those presented here. Große-Brauckmann proved in [Gro93] that each Delaunay surface is the associated MC 1 surface of a helicoid in $\mathbb{S}^3$. The family of Delaunay surfaces consists of embedded (undoloid) and non-embedded (nodoid) examples. In order to construct non-Alexandrov embedded examples one has to consider the minimal surface from Proposition 6.6, and choose the positive oriented surface, i.e. the upward-pointing surface normal ν. If (c_i', η, ν), $i = 0, 2$ is positively oriented, then $\langle c_i', \xi \rangle = -1$. Therefore the twist α is decreasing which implies $\tilde{\kappa} = 1 - \alpha' > 1$ for $\tilde{c}_{0/2}$. With this setup one has to solve the second period problem.

As far as the author knows it is an open problem to show the convergence of the ends of the CMC surface. It would be nice to proof, that M has catenoidal ends, in the sense they are described in [DH09]. To show this one may consider the minimal surface in $\mathrm{Nil}_3(\mathbb{R})$ and wrap it between two copies of horizontal helicoids from [DH09] which differ by a vertical translation about height h:

$$\begin{aligned}
H_1^1(u_1, v_1) &= \frac{\sinh(\alpha v_1)}{\alpha(\psi'(u_1) - \alpha)} \cos\psi(u_1) \\
H_1^2(u_1, v_1) &= -G(u_1) \\
H_1^3(u_1, v_1) &= \frac{-\sinh(\alpha v_1)}{\alpha(\psi'(u_1) - \alpha)} \sin\psi(u_1), \text{ and} \\
H_2^1(u_2, v_2) &= \frac{\sinh(\alpha v_2)}{\alpha(\psi'(u_2) - \alpha)} \cos\psi(u_2) \\
H_2^2(u_2, v_2) &= -G(u_2) \\
H_2^3(u_2, v_2) &= \frac{-\sinh(\alpha v_2)}{\alpha(\psi'(u_2) - \alpha)} \sin\psi(u_2) + h.
\end{aligned}$$

Where α is chosen such that the width of the helicoid is $a \sin\varphi$, which is possible by Lemma 5.1. One has to show that the helicoids are barriers from below and above of the minimal surface $M(a, b_0)$.

The idea is to prove exponential convergence of the two helicoids, i.e. there exists $C, \lambda \in \mathbb{R}$ independent of $x \in H_1$ such that

$$d(x, H_2) < Ce^{-\lambda|x|}, \quad \text{for } |x| \to \infty. \tag{6.5}$$

Since the Plateau solution $M(a, b_0)$ is bounded by H_1 and H_2 for an appropiate choice of h, this proves that $M(a, b_0)$ has a helicoidal end and since the convergence is exponential this translates to the sister surface.

References

[AR05] Uwe Abresch and Harold Rosenberg, *Generalized Hopf differentials*, Mat. Contemp. **28** (2005), 1–28 (English).

[CH12] S. Cartier and L. Hauswirth, *Deformations of constant mean curvature 1/2 surfaces in H2xR with vertical ends at infinity*, ArXiv e-prints (2012).

[CM11] Tobias H. Colding and William P. Minicozzi II, *A course in minimal surfaces*, Providence, RI: American Mathematical Society (AMS), 2011 (English).

[Dan07] Benoît Daniel, *Isometric immersions into 3-dimensional homogeneous manifolds*, Comment. Math. Helv. **82** (2007), no. 1, 87–131 (English).

[DH09] Benoît Daniel and Laurent Hauswirth, *Half-space theorem, embedded minimal annuli and minimal graphs in the Heisenberg group*, Proceedings of the London Mathematical Society **98** (2009), no. 2, 445–470.

[Dou31] Jesse Douglas, *Solution of the problem of Plateau*, Trans. Am. Math. Soc. **33** (1931), 263–321 (English).

[FM07] Isabel Fernández and Pablo Mira, *Harmonic maps and constant mean curvature surfaces in* $\mathbb{H}^2 \times \mathbb{R}$., Am. J. Math. **129** (2007), no. 4, 1145–1181 (English).

[GK10] Karsten Große-Brauckmann and Robert B. Kusner, *Conjugate Plateau constructions for homogeneous 3-manifolds*, preprint, 2010.

[Gro93] Karsten Große-Brauckmann, *New surfaces of constant mean curvature*, Math. Z. **214** (1993), no. 4, 527–565 (English).

[GT01] David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, reprint of the 1998 ed., Berlin: Springer, 2001 (English).

[Gul73] Robert D. Gulliver, *Regularity of minimizing surfaces of prescribed mean curvature*, The Annals of Mathematics **97** (1973), no. 2, pp. 275–305 (English).

[HRS08] Laurent Hauswirth, Harold Rosenberg, and Joel Spruck, *On complete mean curvature* $1/2$ *surfaces in* $\mathbb{H}^2 \times \mathbb{R}$, Commun. Anal. Geom. **16** (2008), no. 5, 989–1005 (English).

[HS88] Joel Hass and Peter Scott, *The existence of least area surfaces in 3-manifolds*, Trans. Am. Math. Soc. **310** (1988), no. 1, 87–114 (English).

[Kar89] Hermann Karcher, *The triply periodic minimal surfaces of Alan Schoen and their constant mean curvature companions.*, Manuscr. Math. **64** (1989), no. 3, 291–357 (English).

[Law70] H. Blaine Jr. Lawson, *Complete minimal surfaces in* $\mathbb{S}^3$, The Annals of Mathematics **92** (1970), no. 3, pp. 335–374 (English).

[Mor66] Charles Bradfield jun. Morrey, *Multiple integrals in the calculus of variations*, Die Grundlehren der mathematischen Wissenschaften, no. 130, Berlin-Heidelberg-New York: Springer-Verlag, 1966 (English).

[MT11] José Miguel Manzano and Francisco Torralbo, *New examples of constant mean curvature surfaces in* $\mathbb{S}^2 \times \mathbb{R}$ *and* $\mathbb{H}^2 \times \mathbb{R}$, arXiv:1104.1259v2 [math.DG], 2011.

[MY82] William H. Meeks III and Shing-Tung Yau, *The existence of embedded minimal surfaces and the problem of uniqueness*, Mathematische Zeitschrift **179** (1982), 151–168, 10.1007/BF01214308.

[Nit75] Johannes C.C. Nitsche, *Vorlesungen über Minimalflächen*, Die Grundlehren der mathematischen Wissenschaften, no. 199, Berlin-Heidelberg-New York: Springer, 1975 (German).

[Oss70] Robert Osserman, *A proof of the regularity everywhere of the classical solution to Plateau's problem*, The Annals of Mathematics **91** (1970), no. 3, pp. 550–569 (English).

[Ple12] Julia Plehnert, *Constant mean curvature surfaces in homogeneous manifolds.*, Logos Verlag, 2012 (English).

[Rad30] Tibor Radó, *The problem of the least area and the problem of Plateau*, Mathematische Zeitschrift **32** (1930), 763–795 (English).

Technische Universität Darmstadt, Fachbereich Mathematik, Geometrie und Approximation, Schlossgartenstr. 7, 64289 Darmstadt, Germany

E-mail address: `plehnert@mathematik.tu-darmstadt.de`